

§ 180.544

40 CFR Ch. I (7–1–11 Edition)

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[65 FR 12134, Mar. 8, 2000]

§ 180.544 Methoxyfenozide; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the insecticide methoxyfenozide per se; benzoic acid, 3-methoxy-2-methyl-, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl) hydrazide in or on the following food commodities:

Commodity	Parts per million
Acerola	0.4
Almond, hulls	25
Animal feed, nongrass, group 18, forage	50.0
Animal feed, nongrass, group 18, hay	150.0
Apple, wet pomace	7.0
Artichoke, globe	3.0
Avocado	0.6
Bean, dry, seed	0.24
Brassica, head and stem, subgroup 5A	7.0
Brassica, leafy greens, subgroup 5B	30
Bushberry subgroup 13-07B	3.0
Canistel	0.6
Cattle, fat	0.50
Cattle, meat	0.02
Coriander, leaves	30
Corn, field, forage	15
Corn, field, grain	0.05
Corn, field, refined oil	0.20
Corn, field, stover	125
Corn, pop, grain	0.05
Corn, pop, stover	125
Corn, sweet, forage	30
Corn, sweet, kernel plus cob with husks removed	0.05
Corn, sweet, stover	60
Cotton, gin byproducts	35
Cotton, undelinted seed	2.0
Cranberry	0.5
Feijoa	0.4
Fruit, pome, group 11	1.5
Fruit, stone, group 12, except plum, prune, fresh	3.0
Goat, fat	0.50
Goat, meat	0.02
Grain, aspirated fractions	2.0
Grape	1.0
Grape, raisin	1.5
Grass, forage, fodder and hay, group 17, forage	18.0
Grass, forage, fodder and hay, group 17, hay ..	30.0
Guava	0.4
Hog, fat	0.1
Hog, meat	0.02
Horse, fat	0.50
Horse, meat	0.02
Jaboticaba	0.4
Leaf petioles subgroup 4B	25
Leafy greens subgroup 4A	30
Longan	2.0
Lychee	2.0
Mango	0.6
Milk	0.10
Nut, tree, group 14	0.10

Commodity	Parts per million
Okra	2.0
Onion, green, subgroup 3-07B	5.0
Papaya	0.6
Passionfruit	0.4
Pea and bean, succulent shelled, subgroup 6B	0.2
Pea, blackeyed, seed	4.0
Pea, dry seed	2.5
Pea, southern, seed	4.0
Peanut	0.02
Peanut, hay	55.0
Peanut, oil	0.04
Peppermint, tops	7.0
Pistachio	0.10
Plum, prune, fresh	0.30
Pomegranate	0.6
Poultry, fat	0.02
Poultry, meat	0.02
Pulasan	2.0
Rambutan	2.0
Sapodilla	0.6
Sapote, black	0.6
Sapote, mamey	0.6
Sheep, fat	0.50
Sheep, meat	0.02
Soybean, aspirated grain fractions	160
Soybean, forage	30
Soybean, hay	80
Soybean, hulls	2.0
Soybean, seed	1.0
Spanish lime	2.0
Spearmint, tops	7.0
Star apple	0.6
Starfruit	0.4
Strawberry	1.5
Turnip, greens	30
Vegetable, cucurbit, group 9	0.3
Vegetable, foliage of legume, except soybean, subgroup 7A	35
Vegetable, fruiting, group 8	2.0
Vegetable, leaves of root and tuber, group 2 ...	30
Vegetable, legume, edible podded, subgroup 6A	1.5
Vegetable, root, subgroup 1A	0.5
Vegetable, tuberous and corm, except potato, subgroup 1D	0.02
Wax jambu	0.4

(2) For combined residues of the insecticide methoxyfenozide; benzoic acid, 3-methoxy-2-methyl-, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl) hydrazide and its glucuronide metabolite RH-141,518; β -D-Glucopyranuronic acid, 3-[2-(1,1-dimethylethyl)-2-(3,5-dimethylbenzoyl)-hydrazino]carbonyl-2-methylphenyl-] in the following commodities:

Commodity	Parts per million
Cattle, liver	0.40
Cattle, meat byproducts, except liver	0.10
Egg	0.02
Goat, liver	0.40
Goat, meat byproducts, except liver	0.10
Hog, liver	0.1
Hog, meat byproducts, except liver	0.02
Horse, liver	0.40
Horse, meat byproducts, except liver	0.10

Environmental Protection Agency

§ 180.546

Commodity	Parts per million
Poultry, liver	0.10
Poultry, meat byproducts, except liver	0.02
Sheep, liver	0.40
Sheep, meat byproducts, except liver	0.10

(b) *Section 18 emergency exemptions.* Time-limited tolerances specified in the following table are established for residues of the insecticide methoxyfenozide *per se* (benzoic acid, 3-methoxy-2-methyl-, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl) hydrazide), in or on the specified agricultural commodities, resulting from use of the pesticide pursuant to FFIFRA section 18 emergency exemptions. The tolerances expire and will be revoked on the date specified in the table.

Commodity	Parts per million	Expiration/revocation date
Sorghum, forage	30.0	12/31/12
Sorghum, grain	0.05	12/31/12
Sorghum, stover	60.0	12/31/12

(c) *Tolerances with regional registrations.* Tolerances are established for residues of the insecticide methoxyfenozide, including its metabolites and degradates. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only methoxyfenozide, benzoic acid, 3-methoxy-2-methyl-, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl) hydrazide.

Commodity	Parts per million
Citrus, Oil	100
Fruit, citrus, group 10	10

(d) *Indirect or inadvertent residues.* [Reserved]

[67 FR 59203, Sept. 20, 2002, as amended at 68 FR 32389, May 30, 2003; 68 FR 37765, June 25, 2003; 69 FR 58097, Sept. 29, 2004; 70 FR 7047, Feb. 10, 2005; 70 FR 51604, Aug. 31, 2005; 70 FR 75739, Dec. 21, 2005; 71 FR 32853, June 7, 2006; 73 FR 11826, Mar. 5, 2008; 74 FR 22468, May 13, 2009; 74 FR 45335, Sept. 2, 2009; 76 FR 34886, June 15, 2011]

§ 180.545 Prallethrin (RS)-2-methyl-4-oxo-3-(2-propynyl)cyclopent-2-enyl (1RS)-cis, trans-chrysanthemate; tolerances for residues.

(a) *General.* (1) A tolerance of 1.0 ppm is established for residues of the insecticide prallethrin (RS)-2-methyl-4-oxo-3-(2-propynyl)cyclopent-2-enyl (1RS)-cis, trans-chrysanthemate as follows:

(2) In or on food commodities in food handling establishments where food and food products are held, processed, prepared and/or served.

(3) Application shall be limited to space, general surface, and spot and/or crack and crevice treatment in food handling establishments where food and food products are held, processed, prepared and/or served. General surface or space spray applications may be used only when the facility is not in operation provided exposed food has been covered or removed from the area being treated prior to application. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application. Spray concentrate shall be limited to a maximum of 2.0% active ingredient. Contamination of food or food contact surfaces shall be avoided. Food contact surfaces and equipment should be thoroughly washed with an effective cleaning compound and rinsed with potable water after use of the product.

(4) To assure safe use of the additive, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency, and it shall be used in accordance with such label and labeling.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[65 FR 39313, June 26, 2000, as amended at 71 FR 74819, Dec. 13, 2006]

§ 180.546 Mefenoxam; tolerances for residues.

(a) *General.* Tolerances are established for residues of mefenoxam, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance